arXiv:1208.0191v1 [hep-ph] 1 Aug 2012

EFFECT OF LEE-WICK THERMODYNAMICS IN THE COSMOLOGY OF THE EARLY UNIVERSE

KAUSHIK BHATTACHARYA

Department of Physics, Indian Institute of Technology, Kanpur,
Kanpur 208016, India,
kaushikb@iitk.ac.in

SURATNA DAS

Tata Institute of Fundamental Research, Homi Bhabha Road,
Colaba, Mumbai 400005, India
suratna@tifr.res.in

The Lee-Wick theories require unusual Lee-Wick (LW) partners to the standard model (SM) particles. The excitations of the unusual fields may have indefinite norms in the Hilbert space. In the present talk the thermodynamic results of a toy LW universe (where each normal particle has only one LW partner and the standard massless gauge fields do not appear in the normal particle spectrum as their LW partners has the potential to produce negative energy density) will be discussed. It will be shown that in the LW universe the time-temperature relation gets modified. This time-temperature relationship shows interesting new physics. More over we see how a LW particle dominated early cosmology transforms into the standard cosmology. A brief discussion on the realistic LW universe containing normal gauge bosons and two fermionic partners for each SM chiral fermion will be presented at the end.

1. The toy model

To tackle the problem of divergences in quantum field theories (QFTs) the LW theories[1] assume a LW partner for each SM field. It turns out that to attack the problem of divergences in QFT one has to quantize the LW partners in such a way that their excitations may have indefinite norms in the Hilbert space. In the present talk we will utilize the LW standard model[2] to investigate the physics of the early universe and we will mainly follow Ref. 3. In the standard model of cosmological evolution the energy density, pressure of ultra-relativistic bosons and fermions are given as

$$\rho_b^{(\rm sm)} = \frac{g\pi^2 T^4}{30}\,,\ \ p_b^{(\rm sm)} = \frac{g\pi^2 T^4}{90}\,, \tag{1}$$

and

$$\rho_f^{(\rm sm)} = \frac{7g\pi^2 T^4}{240}\,,\ \ p_f^{(\rm sm)} = \frac{7g\pi^2 T^4}{720}\,, \tag{2}$$

where the subscripts b and f stands for bosons and fermions and g stands for any internal degree of freedom of the ultra-relativistic species.

In the toy model it is assumed that each SM field is accompanied by a LW partner. In this simplified picture the energy density, pressure for each of these partners are given by:[4,5]

$$\rho_b^{(\mathrm{LW})} = -g\left(\frac{\pi^2 T^4}{30} - \frac{M^2 T^2}{24}\right)\,, \quad p_b^{(\mathrm{LW})} = -g\left(\frac{\pi^2 T^4}{90} - \frac{M^2 T^2}{24}\right)\,, \tag{3}$$

and for the fermionic LW partners:

$$\rho_f^{(\mathrm{LW})} = -g\left(\frac{7\pi^2 T^4}{240} - \frac{M^2 T^2}{48}\right)\,, \quad p_f^{(\mathrm{LW})} = -g\left(\frac{7\pi^2 T^4}{720} - \frac{M^2 T^2}{48}\right)\,, \tag{4}$$

In the above equations M is the mass of a generic LW partner and as the system is relativistic $T > M$. In general the masses of the LW partners are supposed to be much greater than their standard model counterparts. As the LW partner states can have indefinite norm, they are barred to appear as in-states or out-states in any S-matrix process. The LW partners can only appear as resonances in processes involving SM particles. The LW resonances may thermalize (in the cosmic plasma) if the SM particles energies are high i.e., $T > M$. If $T < M$ the LW resonances decouple and turn out to be non-thermal.

If $\tilde{M}$ is the mass of the maximally massive partner field, then the net energy density and entropy density of the relativistic fields in the early universe composed of the SM fields and there LW partners turns out to be,

$$\rho = \frac{\tilde{M}^2}{24}\tilde{g}_* T^2\,, \tag{5}$$

where $\tilde{g}_*$ is the effective degree of freedom for energy calculation, and

$$s = \frac{\tilde{M}^2}{12}\tilde{g}_{*s} T \tag{6}$$

where $\tilde{g}_{*s}$ is the effective degree of freedom for entropy calculation.

The effective degrees of freedom are as

$$\tilde{g}_* = \sum_{i=\mathrm{bosons}} g_i \left(\frac{M_i}{\tilde{M}}\right)^2 \left(\frac{T_i}{T}\right)^2 + \frac{1}{2}\sum_{i=\mathrm{fermions}} g_i \left(\frac{M_i}{\tilde{M}}\right)^2 \left(\frac{T_i}{T}\right)^2\,, \tag{7}$$

and

$$\tilde{g}_{*s} = \sum_{i=\mathrm{bosons}} g_i \left(\frac{M_i}{\tilde{M}}\right)^2 \left(\frac{T_i}{T}\right) + \frac{1}{2}\sum_{i=\mathrm{fermions}} g_i \left(\frac{M_i}{\tilde{M}}\right)^2 \left(\frac{T_i}{T}\right)\,. \tag{8}$$

Here it has been assumed that each SM boson (or fermion) and its LW partner are in thermal equilibrium but all the different species of particles and their partners may not be sharing the same temperature. In the above relations $\tilde{M}$ is the mass of the maximally massive LW partner which is relativistic at temperature T. The above equations for energy density and entropy density are correct if the spectrum of the particles does not contain any massless gauge bosons. In presence of massless gauge bosons, in the realistic model, the above relations get modified.

Using the above forms of energy density and entropy density one can show the time-temperature relation for the LW universe to be:

$$T_{\mathrm{GeV}} \sim 10^{-24}(\pi\tilde{g}_*)^{-1/2}\left(\frac{M_{\mathrm{Pl}}}{\tilde{M}}\right)\frac{1}{t_{\mathrm{sec}}}\,, \tag{9}$$

The time-temperature relation is plotted in Fig. 1. The bump appears at a specific time in Fig. 1 because at that time the heaviest LW partner with mass $\tilde{M}$ becomes non-relativistic and the the heaviest mass value in Eq. (9) is replaced by the next heaviest LW partner mass $\tilde{M}'$. As soon this happens the temperature of the system shoots up ($T \propto 1/\tilde{M}$ and $\tilde{M} > \tilde{M}'$) and the previous massive LW partner re-thermalize. This happens whenever the new temperature, T', is such that $T' > \tilde{M}$. Then again the temperature starts to fall up to a point beyond which the new temperature shoots up and the previous LW partner re-thermalize. This cycle repeats for some time until the temperature rise due to the decoupling of the heaviest LW partner falls below $\tilde{M}$. This is an out of equilibrium process as the entropy also shoots up during this time as shown in Fig. 2.

As the LW partners become non-relativistic they decouple from the plasma and do not contribute in the energy density or entropy density of the system. After some of the LW partners decouple, the modified entropy density and energy density of the cosmic plasma becomes

$$\rho' = \frac{\tilde{M'}^2}{24}\tilde{g'}_* T'^2 + \frac{\pi^2 T'^4}{30}\, g'_*\,, \tag{10}$$

$$s' = \frac{\tilde{M'}^2}{12}\tilde{g'}_{*s} T' + \frac{2\pi^2 T'^3}{45}\, g'_{*s}\,. \tag{11}$$



Fig. 1. Figure showing the bump in the temperature as the heaviest Lee-Wick partner with mass 10^9 GeV becomes non-thermal and its position is taken up by the next heaviest Lee-Wick partner of mass 3×10^8 GeV. For a simple illustration the effective degrees of freedom are assumed to remain constant in the process, $\tilde{g}_* \sim 80$. The green region in the plot represents a region of rapid and violent temperature fluctuations where the system has gone out of equilibrium.

The first terms on the right hand sides of the equations comes from that part of the plasma which consists of SM particles and their LW partners which are still thermalized at that temperature T'. The second term of the right hand sides of the equations comes from a plasma constituted by only SM particles whose LW partners have already decoupled from the cosmic plasma. As the LW partners are heavier than their SM partners it is safe to assume that these SM particles are still relativistic at temperature T'. In the above equations we assume that $\tilde{M}'$ is the mass of the maximally massive Lee-Wick partner at the corresponding temperature T'. The effective degrees of freedom changes to $\tilde{g'}_*$ and $\tilde{g'}_{*s}$ from their previous values. Also, g'_* and g'_{*s} account for the effective degrees of freedom of those SM particles whose Lee-Wick partners have decoupled. It may happen that after some decouplings of the LW partners the energy density and entropy density transforms to the standard cosmological forms because the T'^4 term and T'^3 terms start to dominate over the other terms.

2. The realistic model

In the realistic model there are two fermionic partners of a standard chiral fermion and there is one more degree of freedom for the massive partners of the massless



Fig. 2. Figure showing the bump in the entropy density of the universe as the heaviest Lee-Wick partner with mass 10^9 GeV becomes non-thermal and its position is taken up by the next heaviest Lee-Wick partner of mass 3×10^8 GeV. The effective degrees of freedom are assumed to remain constant in the process as $\tilde{g}_{*s} \sim 80$. The SM particle freed of its partner is assumed to be a boson with $g = 2$. The spiked green region in the plot represents a region of rapid and violent entropy fluctuations where the system has gone out of equilibrium.

gauge bosons. In this case:

$$\rho = \frac{\tilde{M}^2}{24}\tilde{g}_{*N}T^2 - \frac{7\pi^2}{240}\tilde{g}_F T^4 - \frac{\pi^2}{30}nT^4\,, \tag{12}$$

where the new degrees of freedom $\tilde{g}_{*N}$ originates from the standard bosonic contribution, similar to the bosonic contribution in Eq. (7). As there are more LW fermions, the unpaired fermionic contribution comes with $\tilde{g}_F$. The last term in Eq. (12) arises from the extra longitudinal degree of freedom of the massive LW partners of the normal massless gauge bosons. Here n denotes the number of massive vector boson partners of SM gauge bosons. In the last equation it is assumed that all the species in the plasma are in thermal equilibrium.

In the present scenario one sees that if there are two LW fermionic partners for each standard fermion and the longitudinal mode of the LW partners of the normal gauge bosons also contribute to the energy density then the net energy density is not positive definite. There can be two ways in which one can obtain a positive definite energy density of the early universe infested with LW partners. The first one depends upon the principle of parametric resonance during preheating after inflation.[6] During these phase only massive bosons are produced and if these massive bosons initially thermalize during reheating then the problem with the fermions and gauge bosons disappear. The second option is related to the reheating temperature of the universe. If the LW fermion partners and the LW partners of the gauge bosons decoupling temperature is more than the reheat temperature then both of these species will not be present in the energy density term and henceforth the energy density will be positive definite. These scenario depends upon the mass hierarchy of the LW partners.

In conclusion one can say that a realistic LW radiation dominated universe is a difficult thing to realize but for certain circumstances as discussed above one can indeed have a LW radiation phase during/after reheating.

References

1. T. D. Lee and G. C. Wick, Phys. Rev. D **2**, 1033 (1970).
2. B. Grinstein, D. O'Connell and M. B. Wise, Phys. Rev. D **77**, 025012 (2008) [arXiv:0704.1845 [hep-ph]].
3. K. Bhattacharya and S. Das, Phys. Rev. D **86**, 025009 (2012) [arXiv:1203.1109 [hep-ph]].
4. B. Fornal, B. Grinstein and M. B. Wise, Phys. Lett. B **674**, 330 (2009) [arXiv:0902.1585 [hep-th]].
5. K. Bhattacharya and S. Das, Phys. Rev. D **84**, 045023 (2011) [arXiv:1108.0483 [hep-ph]].
6. L. Kofman, A. D. Linde and A. A. Starobinsky, Phys. Rev. Lett. **73**, 3195 (1994) [hep-th/9405187].